


# Continuously cored topsets of the Albian–Cenomanian(?) Nanushuk Formation, Wainwright No. 1 well, northwestern Alaska: Deposition in a marine-influenced delta plain setting

David L. LePain<sup>1</sup> and Paul L. Decker<sup>2</sup>  
<sup>1</sup> Alaska Division of Geological & Geophysical Surveys, 3354 College Road, Fairbanks, AK 99709  
<sup>2</sup> Alaska Division of Oil and Gas, 555 West 7th Avenue, Suite 800, Anchorage, AK 99501



Explanation		Facies Codes	
21.7, 151 ▶ Porosity (percent), permeability (md)	Trough or planar cross-stratification	FI - laminated mudstone	
Sandstone	Micro-extensional faults	Fm - massive mudstone	
Mudstone	Hummocky cross-stratification	Fc - carbonaceous mudstone	
Carbonaceous mudstone	Plane-parallel lamination	Fc - carbonaceous mudstone	
Coal	Convolute lamination/bedding	C - coal	
Gradational change in lithology	Rosselia	Sr - ripple cross-laminated sandstone	Sm - massive sandstone
Gradational change in facies	Bioturbation	Sx - cross-bedded sandstone	Scb - convolute bedded sandstone
Current ripple cross-lamination	Rhizolith (roots)	Sif - plane-parallel laminated fine-grained sandstone	Sb - burrow-mottled sandstone
Wave ripple cross-lamination	Syneresis cracks	Sfm - plane-parallel laminated medium-grained sandstone	Sfh - heterolithic sandstone and mudstone
	Breccia-like fabric		

Grain size explanation	
Ivf - lower very fine sand	
If - lower fine sand	
Im - lower medium sand	

*Bioturbation Index from Droser and Botzjer, 1986*